

Precision measurement of the η -meson mass at COSY-ANKE

Paul Goslawski^(a), Alfons Khoukaz^(a), Malte Mielke^(a), Michael Papenbrock^(a), Daniel Schroeer^(a), Alexander Täschner^(a) for the ANKE collaboration

^(a) Institut für Kernphysik, Universität Münster, D-48149 Münster, Germany

Measurements of the mass of the η -meson performed at different experimental facilities (i.e., COSY-GEM, MAMI, CLEO, KLOE, NA48) over the last decade have resulted in very precise data which differ by up to $0.5 \text{ MeV}/c^2$, i.e., more than eight standard deviations. In order to clarify this situation a new measurement of the $dp \rightarrow {}^3\text{He}\eta$ reaction near threshold was proposed at the COoler SYnchrotron - COSY - of the Forschungszentrum Jülich with the aim to achieve a mass resolution of $\Delta m < 50 \text{ keV}/c^2$.

Through the measurement of a set of twelve beam momenta p_d and associated ${}^3\text{He}\eta$ final state momenta p_f a value for the η mass can be obtained on the basis of pure kinematics by the determination of the production threshold. The individual beam momenta were fixed with a relative precision of better than 10^{-4} at $3 \text{ GeV}/c$ by using a polarized deuteron beam and inducing an artificial depolarizing resonance, which occurs at a well-defined frequency. The final-state momenta in the two-body $dp \rightarrow {}^3\text{He}\eta$ reaction were investigated in detail by studying the size of the ${}^3\text{He}$ momentum ellipse in the focal plane of the ANKE spectrometer. The method for determination of the η mass as well as final results will be discussed in this presentation.

Supported by the COSY-FFE grants.

E-mail: paul.goslawski@uni-muenster.de