

From transition form factors to $(g - 2)_\mu$

Bastian Kubis¹, Michael C. Kunkel²

¹Helmholtz-Institut für Strahlen- und Kernphysik, Universität Bonn

²Institut für Kernphysik (IKP-1), Forschungszentrum Jülich

Hadron Physics Summer School 2016

The charge distributions of composite objects are characterized by *form factors*. Whether describing the cross sections for electron scattering off nucleons (composed of quarks and gluons) or nuclei (composed of nucleons), form factors parametrize the deviation from the scattering amplitude off a point-like object (like an electron); to study them allows us to learn something about the constituents as well as the strong interactions that bind them together [1]. The momentum dependence of the form factors can therein be understood as the Fourier transform of the charge distribution (within a certain reference frame).

Similar form factors can also be used to characterize the electromagnetic *transition* between different particles. In particular, we can also investigate the form factors of particles that cannot be prepared as targets in scattering experiments through the study of their *decays* into final states including an electron–positron pair. Of particular recent interest are the transition form factors describing the decays of light, flavor-neutral mesons (π^0 , η , η') into one real and one virtual photon; with only one single hadron involved, it is this hadron’s electromagnetic structure that can be understood in such decays.

In this working group, we want to concentrate on the decay $\eta' \rightarrow \gamma e^+ e^-$. The decaying η' can be produced in the reaction $e^- p \rightarrow e^- \eta' p$ (electroproduction), for instance at the CLAS experiment at Jefferson Lab (Virginia, USA). We want to understand the existing experimental results for such form factors [2], how they can be understood and interpreted theoretically [3], and what their implications are for a wider range of modern particle physics problems of current interest [4]. Finally, we want to work on a proposal for improved experiments to measure the transition form factors.

References

- [1] *see e.g.:* F. Halzen and A. D. Martin, *Quarks And Leptons: An Introductory Course In Modern Particle Physics*, New York, USA: Wiley (1984).
- [2] M. Ablikim *et al.* [BESIII Collaboration], *Observation of the Dalitz Decay $\eta' \rightarrow \gamma e^+ e^-$* , Phys. Rev. D **92** (2015) 012001 [arXiv:1504.06016 [hep-ex]].
- [3] C. Hanhart, A. Kupść, U.-G. Meißner, F. Stollenwerk and A. Wirzba, *Dispersive analysis for $\eta \rightarrow \gamma \gamma^*$* , Eur. Phys. J. C **73** (2013) 2668 [arXiv:1307.5654 [hep-ph]].
- [4] E. Czerwiński, S. Eidelman, C. Hanhart, B. Kubis, A. Kupść, S. Leupold, P. Moskal and S. Schandmand, *MesonNet Workshop on Meson Transition Form Factors*, arXiv:1207.6556 [hep-ph].