


EDMs of Light Nuclei

Jülich-Bonn Collaboration (JBC):

J. Bsaisou, C.Hanhart, S.Liebig, U.-G.Meißner, D.Minossi,
A.Nogga, J.de Vries, A.Wirzba

09.10.2013 | Jan Bsaisou

EDMs from \mathcal{CP} Formfactor (F_3):



Outline:

- **CP-violation** beyond CKM matrix in the SM: \mathcal{L}_{QCD} θ -term (dim. 4)
 - EDM of the deuteron / EDM of helium-3
 - strategies of testing whether $\bar{\theta}$ -term is the origin
- **CP-violation** from physics beyond the SM: SUSY, multi-Higgs, ...
 - dim. 6 sources: qEDM, qCEDM, gCEDM, 4qEDMs
 - EDM of the deuteron / EDM of helium-3
 - disentangling dim. 6 sources

The \mathcal{L}_{QCD} θ -Term

topologically non trivial vacuum \rightarrow \mathcal{CP} term in \mathcal{L}_{QCD} :

$$\mathcal{L} = \mathcal{L}_{QCD}^{CP} + \theta \frac{g_S^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a,\mu\nu}$$

$$\dots + \theta \frac{g_S^2}{32\pi^2} G_{\mu\nu}^a \tilde{G}^{a,\mu\nu} \xrightarrow{U_A(1)} \dots - \bar{\theta} m^* \sum_f \bar{q}_f i\gamma_5 q_f$$

with $\bar{\theta} = \theta + \arg \det \mathcal{M}$, naive dim. analysis (NDA): $\bar{\theta} \sim \mathcal{O}(1)$


\mathcal{M} : quark mass matrix, $m^* = \frac{m_u m_d}{m_u + m_d}$

Physics Beyond SM (BSM):

SUSY, multi-Higgs, Left-Right-Symmetric models, ...

Effective field theory approach:


- All degrees of freedom beyond a specified scale are integrated out:
 ↪ Only SM degrees of freedom remain: q, g, H, W^\pm, \dots
- Relics of eliminated BSM physics ‘remembered’ by the values of the **low-energy constants (LECs)** of the **CP-violating contact terms**, e.g.



BSM physics continued: CP-violating dim. 6 sources

Removal of the Higgs and transition to hadronic fields (plus mixing)

Add to SM all possible T- and P-odd contact interactions






θ -Term on the Hadronic Level

hadronic level: non perturbative techniques required: e.g. 2-flavor *ChPT*


- Symmetries of QCD preserved by the effective field theory (EFT)

$$\mathcal{L}_{\text{QCD}}^\theta = -\bar{\theta} m^* \sum_f \bar{q}_f i \gamma_5 q_f: \quad \mathcal{CP}, \text{I} \quad \Leftrightarrow \quad \mathcal{M} \rightarrow \mathcal{M} + \bar{\theta} m^* i \gamma_5 \quad m^* = \frac{m_u m_d}{m_u + m_d}$$

	\mathcal{CP}, I	\mathcal{CP}, I						
				$\mathcal{CP}, \text{I} + \text{I}$				
$\mathcal{L}_\theta^{\text{ChPT}} =$	$g_0^\theta N^\dagger \vec{\pi} \cdot \vec{\tau} N$	+	$g_1^\theta N^\dagger \pi_3 N$	+	$N^\dagger (b_0 + b_1 \tau_3) S^\mu \nu^\nu F_{\mu\nu} N$	+	\dots	
	\Downarrow		\Downarrow		\Downarrow			
	$\begin{array}{c} \\ \pi^\pm, \pi^0 \\ \\ N \end{array}$		$\begin{array}{c} \\ \pi^0 \\ \\ N \end{array}$		$\begin{array}{c} \\ \gamma \\ \\ N \end{array}$		\dots	
								
	dominating for ${}^3\text{He}$		dominating for D		important for p, n			

Lebedev et al. (2004), Mereghetti et al. (2010), J.B. et al. (2013)

Selection of the Ground State: θ -term



ground state fixed under \mathcal{CP} : readjustment of coordinates
 \rightarrow no pion tadpole

\Rightarrow impact on πN -sector:

$$\mathcal{L}_{\pi N} = \underbrace{g_0^\theta N^\dagger \vec{\pi} \cdot \vec{\tau} N}_{\mathcal{CP}, I} + \dots \longrightarrow \underbrace{(g_0^\theta + \delta g_0^\theta) N^\dagger \vec{\pi} \cdot \vec{\tau} N}_{\mathcal{CP}, I} + \underbrace{g_1^\theta N^\dagger \pi_3 N}_{\mathcal{CP}, I} + \dots$$

$$\delta g_0^\theta = \mathcal{O}((\delta M_\pi^2)_{QCD})$$

$$(\delta M_\pi^2)_{QCD} = (M_{\pi^+}^2 - M_{\pi^0}^2)_{QCD}$$

θ -term: \mathcal{CP} πNN -terms related to LECs c_5 and c_1 :

Crewther et al. (1979); Otnad et al. (2010); Mereghetti et al. (2011);
de Vries et al. (2011); J.B. et al. (2013)

coupling constants g_0^θ, g_1^θ of \mathcal{CP} πNN -vertices can be fixed!

g_0^θ :

$$\mathcal{L}_{\pi N} = \dots + c_5 2B N^\dagger \left((m_u - m_d) \tau_3 + \frac{2m^* \bar{\theta}}{F_\pi} \vec{\pi} \cdot \vec{\tau} \right) N + \dots$$

$$\delta M_{np}^{str} = 4B(m_u - m_d)c_5 \quad \rightarrow \quad g_0^\theta = \bar{\theta} \delta M_{np}^{str} \frac{(1 - \epsilon^2)}{\epsilon} \frac{1}{4F_\pi}$$

$$\delta M_{np}^{em} \quad \rightarrow \quad \delta M_{np}^{str} = (2.6 \pm 0.5) \text{MeV} \quad \text{Walker-Loud et al. (2012)}$$

$$\rightarrow g_0^\theta = (-0.018 \pm 0.007) \bar{\theta}$$

$$\epsilon = (m_u - m_d)/(m_u + m_d), \quad 4Bm^* = M_\pi^2(1 - \epsilon^2), \quad m^* = \frac{m_u m_d}{m_u + m_d}$$

θ -term: \mathcal{CP} πNN -terms related to LECs c_5 and c_1 :

g_1^θ : $\pi_3 NN$ -vertex


$$\epsilon = (m_u - m_d)/(m_u + m_d)$$

$$\mathcal{L}_{\pi N} = \dots + c_1 4B N^\dagger \left((m_u + m_d) + \frac{(\delta M_\pi^2)_{QCD} (1 - \epsilon^2) \bar{\theta}}{2BF_\pi \epsilon} \pi_3 \right) N + \dots$$

1 $c_1 \longleftrightarrow \sigma_{\pi N}$: $c_1 = (-1.0 \pm 0.3) \text{ GeV}^{-1}$

Compilation: Baru et al. (2011)

2 $(\delta M_\pi^2)_{QCD} = \frac{\epsilon^2}{4} \frac{M_\pi^4}{M_K^2 - M_\pi^2}$



$$\longrightarrow g_1^\theta = (0.003 \pm 0.001) \bar{\theta}$$

J.B. et al. (2013)

$$\frac{g_1^\theta}{g_0^\theta} = -0.20 \pm 0.13 \sim \frac{M_\pi}{m_N}$$

$$\gg \epsilon \frac{M_\pi^2}{m_N^2} \sim -0.01 \quad (\text{NDA})$$

$g_0^\theta (\delta M_{np}^{str})$ is unnaturally small

BSM ~~CP~~ sources on the hadronic level

Reliance on Naive Dimensional Analysis (NDA), lattice, ...



$g_0: \mathcal{CP}, I$

$g_1: \mathcal{CP}, I$

$d_0, d_1: \mathcal{CP}, I + I$

$C_{3\pi}: \mathcal{CP}, I$

~~\mathcal{CP}~~
EFT:



θ -term:

$\mathcal{O}(1)$

$\mathcal{O}(M_\pi/m_N)$

$\mathcal{O}(M_\pi^2/m_N^2)$

$\mathcal{O}(\epsilon M_\pi^2/m_N^2)$

qEDM:

$\mathcal{O}(\alpha_{EM}/(4\pi))$

$\mathcal{O}(\alpha_{EM}/(4\pi))$

$\mathcal{O}(1)$

$\mathcal{O}(\alpha_{EM}/(4\pi))$

qCEDM:

$\mathcal{O}(1)$

$\mathcal{O}(1)$

$\mathcal{O}(M_\pi^2/m_N^2)$

$\mathcal{O}(\epsilon M_\pi^2/m_N^2)$

4qLR:

$\mathcal{O}(M_\pi^2/m_N^2)$

$\mathcal{O}(1)$

$\mathcal{O}(M_\pi^2/m_N^2)$

$\mathcal{O}(1)$

gCEDM:

$\mathcal{O}(M_\pi^2/m_N^2)^*$

$\mathcal{O}(M_\pi^2/m_N^2)^*$

$\mathcal{O}(1)$

$\mathcal{O}(\epsilon M_\pi^2/m_N^2)$

4q:

$\mathcal{O}(M_\pi^2/m_N^2)^*$

$\mathcal{O}(M_\pi^2/m_N^2)^*$

$\mathcal{O}(1)$

$\mathcal{O}(\epsilon M_\pi^2/m_N^2)$

*: Goldstone theorem \rightarrow relative $\mathcal{O}(M_\pi^2/m_N^2)$ suppression of $N\pi$ interactions

Nucleon EDM: θ -term case


single nucleon EDM:



“controlled”

→ lattice QCD required

two nucleon EDM:



controlled


isovector

\approx

\ll

isoscalar

Ottnad et al. (2010)



two counter terms

Guo, Meißner (2012)


Sushkov, Flambaum, Khriplovich (1984)

\gg



unknown

D \mathcal{CP} form factor computation technique:




$$G = G_0 + G_0 t_{12} G_0$$

$$t_{12} = (1 - V_{12} G_0)^{-1} V_{12}$$

Note:

- Complementary Monte Carlo based test for plane wave contribution
- Additional analytic computation utilizing PEST separable potential

EDM of the Deuteron at LO: θ -term



LO: ~~$g_0^{\theta} N^{\dagger} \vec{\pi} \cdot \vec{\tau} N (\mathcal{CP}, I)$~~ \rightarrow Isospin select.

NLO: $g_1^{\theta} N^{\dagger} \pi_3 N (\mathcal{CP}, I) \rightarrow$ LO


in units of $g_1^{\theta} e \cdot \text{fm} \cdot (g_A m_N / F_{\pi})$

refs.	potential	no 3P_1 -int	with 3P_1 -int	total
JBC (2013)*	A_{V18}	-1.93×10^{-2}	0.48×10^{-2}	-1.45×10^{-2}
JBC (2013)	CD BONN	-1.95×10^{-2}	0.51×10^{-2}	-1.45×10^{-2}
JBC (2013)*	ChPT(N^2LO) [†]	-1.94×10^{-2}	0.65×10^{-2}	-1.29×10^{-2}
Song (2013)	A_{V18}	-	-	-1.45×10^{-2}
Liu (2004)	A_{V18}	-	-	-1.43×10^{-2}
Afnan (2010)	Reid93	-1.93×10^{-2}	0.40×10^{-2}	-1.43×10^{-2}


*: in preparation †: cutoffs at 600 MeV (LS) and 700 MeV (SFR)

BSM \mathcal{CP} sources: LO $g_1^{\theta} \pi NN$ -vertex also for qCEDM and 4qLR

EDM of the Deuteron: NLO - and N^2LO -Potentials




EDM of the Deuteron: NLO - and N^2LO -Potentials




- ✗: vanishing by selection rules, ✗: sum of diagrams vanishes
✗: vertex correction

EDM of the Deuteron: NLO - and N^2LO -Currents



*: de Vries et al. (2011), J.B. et al. (2013)

EDM of the Deuteron: NLO - and N^2LO -Currents



*: de Vries et al. (2011), J.B. et al. (2013)

- ~~X~~: vanishing by selection rules, ~~X~~: sum of diagrams vanishes

Deuteron EDM from the θ -term

J.B. et al. (2013)




total deuteron EDM: $d_D = d_n + d_p + d_D(2N)$

- single-nucleon contribution: EFT *alone* has no predictive power
 → *Experiment or Lattice QCD* needed in addition
- two-nucleon contribution $d_D(2N)$: EFT *has* predictive power

$$d_D(2N) = \underbrace{-(0.59 \pm 0.39) \cdot 10^{-16} \bar{\theta} \text{ e cm}}_{\text{LO}} + \underbrace{(0.05 \pm 0.02) \cdot 10^{-16} \bar{\theta} \text{ e cm}}_{\text{N}^2\text{LO}}$$


^3He EDM: Power Counting

Utilizing Schroedinger equation $|\psi\rangle = G_0 V|\psi\rangle$ to compare $NN \sim 3N$ ops.


	θ -term	qCEDM	4qLR
	A_θ (LO)	A_{qC} (LO)	A_{4q} ($N^2\text{LO}$)
	$A_\theta \times \frac{M_\pi}{m_N}$ (NLO)	A_{qC} (LO)	$A_{4q} \times \frac{m_N^2}{M_\pi^2}$ (LO)
	$A_\theta \times \frac{M_\pi^2}{m_N^2}$ ($N^2\text{LO}$)	$A_{qC} \times \frac{M_\pi^2}{m_N^2}$ ($N^2\text{LO}$)	$A_{4q} \times \frac{m_N^2}{M_\pi^2}$ (LO)

${}^3\text{He}$ \mathcal{CP} form factor computation technique: Faddeev approach

plane wave:



multiple rescatterings:



$$\langle {}^3\text{He} | V_{\mathcal{CP}}^{12} G_0 \mathcal{O}_3(\vec{q}) | {}^3\text{He} \rangle + \langle {}^3\text{He} | V_{\mathcal{CP}}^{12} G_0 V G \mathcal{O}_3(\vec{q}) | {}^3\text{He} \rangle + \text{perm.}$$


$$\Rightarrow \text{Faddeev equation: } (1 + P) | U_{(3)} \rangle \equiv V G (1 + P) \mathcal{O}_3(\vec{q}) | {}^3\text{He} \rangle$$

$$| U_{(3)}(\vec{q}) \rangle = t_{12} G_0 (1 + P) \mathcal{O}_3(\vec{q}) | {}^3\text{He} \rangle + t_{12} G_0 P | U_{(3)} \rangle + (3NF)$$

$$P = P_{12} P_{23} + P_{13} P_{23}$$

Note: complementary Monte-Carlo based test for plane wave contribution

^3He EDM: quantitative results for g_0 exchange



$$g_0 N^\dagger \vec{\pi} \cdot \vec{\tau} N \quad (\mathcal{CP}, I)$$

$$\theta\text{-term, qCEDM} \quad \rightarrow \quad \text{LO}$$

$$4\text{qLR} \quad \rightarrow \quad \text{N}^2\text{LO}$$

units: $g_0 (g_A m_N / F_\pi) e \text{ fm}$


author	potential	no int.	with int.	total
JBC (2013)*	$A_{V_{18}}\text{UIX}$	-0.45×10^{-2}	-0.13×10^{-2}	-0.57×10^{-2}
JBC (2013)*	CD BONN TM	-0.56×10^{-2}	-0.12×10^{-2}	-0.67×10^{-2}
JBC (2013)*	ChPT (N^2LO) [†]	-0.56×10^{-2}	-0.19×10^{-2}	-0.76×10^{-2}
Song (2013)	$A_{V_{18}}\text{UIX}$	-	-	-0.59×10^{-2}
Stetcu (2008)	$A_{V_{18}}\text{UIX}$	-	-	-1.21×10^{-2}

*: in preparation †: cutoffs at 600 MeV (LS) and 700 MeV (SFR)

Results for ^3H also available (not shown)

JBC (2013)* ~ Song (2013) ~ Stetcu (2008)/2

^3He EDM: quantitative results for g_1 exchange



$$g_1 N^\dagger \pi_3 N \quad (\mathcal{CP}, I)$$

θ -term \rightarrow NLO

qCEDM, 4qLR \rightarrow LO !

units: $g_1 (g_{AMN}/F_\pi) \text{ e fm}$


Ref.	potential	no int.	with int.	total
JBC (2013)*	$A_{V_{18}}\text{UIX}$	-1.09×10^{-2}	-0.02×10^{-2}	-1.11×10^{-2}
JBC (2013)*	CD BONN TM	-1.11×10^{-2}	-0.03×10^{-2}	-1.14×10^{-2}
JBC (2013)*	ChPT ($N^2\text{LO}$) [†]	-1.09×10^{-2}	-0.14×10^{-2}	-0.96×10^{-2}
Song (2013)	$A_{V_{18}}\text{UIX}$	-	-	-1.08×10^{-2}
Stetcu (2008)	$A_{V_{18}}\text{UIX}$	-	-	-2.20×10^{-2}

*: in preparation †: cutoffs at 600 MeV (LS) and 700 MeV (SFR)

Results for ^3H also available (not shown)

JBC (2013)* \sim Song (2013) \sim Stetcu (2008)/2

For completeness: irrelevant results for g_2 exchange



$$g_2 N^\dagger (3 \tau_3 \pi_3 - \vec{\tau} \cdot \vec{\pi}) N \quad (\mathcal{CP}, I)$$

units: $g_2 (g_{AMN}/F_\pi) \text{ e fm}$

Ref.	potential	no int.	with int.	total
JBC (2013)*	$A_{V_{18}} \text{UIX}$	-1.36×10^{-2}	-0.35×10^{-2}	-1.71×10^{-2}
JBC (2013)*	CD BONN TM	-1.46×10^{-2}	-0.37×10^{-2}	-1.83×10^{-2}
JBC (2013)*	ChPT ($N^2\text{LO}$) [†]	-1.42×10^{-2}	-0.14×10^{-2}	-1.56×10^{-2}
Song (2013)	$A_{V_{18}} \text{UIX}$	-	-	-0.66×10^{-2}
Stetcu (2008)	$A_{V_{18}} \text{UIX}$	-	-	-3.40×10^{-2}
Stetcu (2008)	CD BONN TM	-	-	-3.50×10^{-2}

*: in preparation †: cutoffs at 600 MeV (LS) and 700 MeV (SFR)

Results for ${}^3\text{H}$ also available (not shown)

Pattern reinforced: JBC (2013)* \sim Stetcu (2008)/2

Quantitative EDM results in the θ -term scenario

Single Nucleon (with adjusted signs for consistency; note here $e < 0$):

$$\begin{aligned}
 -d_1^{\text{loop}} &\equiv \frac{1}{2}(d_n - d_p)^{\text{loop}} \\
 &= (2.1 \pm 0.9) \cdot 10^{-16} \bar{\theta} \text{ e cm} && \text{(Bsaisou et al. (2013))} \\
 d_n &= +(2.9 \pm 0.9) \cdot 10^{-16} \bar{\theta} \text{ e cm} && \text{(Guo \& Meißner (2012))} \\
 d_p &= -(1.1 \pm 1.1) \cdot 10^{-16} \bar{\theta} \text{ e cm} && \text{(Guo \& Meißner (2012))}
 \end{aligned}$$

Deuteron:

$$\begin{aligned}
 d_D &= d_n + d_p - [(0.59 \pm 0.39) - (0.05 \pm 0.02)] \cdot 10^{-16} \bar{\theta} \text{ e cm} \\
 &= d_n + d_p - (0.54 \pm 0.39) \cdot 10^{-16} \bar{\theta} \text{ e cm} && \text{(Bsaisou et al. (2013))}
 \end{aligned}$$

Helium-3:

$$\begin{aligned}
 d_{^3\text{He}} &= \tilde{d}_n + [(1.52 \pm 0.60) - (0.46 \pm 0.30)] \cdot 10^{-16} \bar{\theta} \text{ e cm} \\
 &= \tilde{d}_n + (1.06 \pm 0.67) \cdot 10^{-16} \bar{\theta} \text{ e cm} && \text{(JBC (2013))}
 \end{aligned}$$

$$\text{with } \tilde{d}_n = 0.88d_n - 0.047d_p \quad \text{(de Vries et al. (2011))}$$

Testing Strategies in the θ EDM scenario

Remember:


$$d_D = d_n + d_p - (0.54 \pm 0.39) \cdot 10^{-16} \bar{\theta} \text{ e cm} \quad (\text{Bsaisou et al. (2013)})$$

$$d_{^3\text{He}} = \tilde{d}_n + (1.06 \pm 0.67) \cdot 10^{-16} \bar{\theta} \text{ e cm} \quad (\text{JBC (2013)})$$

Testing strategies:

- plan A: measure d_n , d_p , and $d_D \xrightarrow{d_D(2N)} \bar{\theta} \xrightarrow{\text{test}} d_{^3\text{He}}$
- plan A': measure d_n , (d_p), and $d_{^3\text{He}} \xrightarrow{d_{^3\text{He}}(2N)} \bar{\theta} \xrightarrow{\text{test}} d_D$
- plan B: measure d_n (or d_p) + Lattice QCD $\rightsquigarrow \bar{\theta} \xrightarrow{\text{test}} d_D$
- plan B': measure d_n (or d_p) + Lattice QCD $\rightsquigarrow \bar{\theta} \xrightarrow{\text{test}} d_p$ (or d_n)


If $\bar{\theta}$ -term tests fail: effective BSM dim. 6 sources: de Vries et al. (2011)



$qEDM$

$$d_D \approx d_p + d_n$$

$$d_{^3He} \approx d_n$$



$qCEDM$

$$d_D > d_p + d_n$$


$$d_{^3He} > d_n$$



$4qLR$

$$d_D > d_p + d_n$$

$$d_{^3He} > d_n$$



$gCEDM + 4qEDM$

$$d_D \sim d_p + d_n$$

$$d_{^3He} \sim d_n$$

→ $g_0, g_1 \propto \alpha/(4\pi)$

$2N$ contribution suppressed by photon loop!

here: only absolute values considered


If $\bar{\theta}$ -term tests fail: effective BSM dim. 6 sources: de Vries et al. (2011)



$qEDM$

$$d_D \approx d_p + d_n$$

$$d_{3He} \approx d_n$$



$qCEDM$

$$d_D > d_p + d_n$$


$$d_{3He} > d_n$$



$4qLR$

$$d_D > d_p + d_n$$

$$d_{3He} > d_n$$



$gCEDM + 4qEDM$

$$d_D \sim d_p + d_n$$

$$d_{3He} \sim d_n$$

→ g_0, g_1

$2N$ contribution enhanced!

here: only absolute values considered


If $\bar{\theta}$ -term tests fail: effective BSM dim. 6 sources: de Vries et al. (2011)



$qEDM$

$$d_D \approx d_p + d_n$$


$$d_{3He} \approx d_n$$



$qCEDM$

$$d_D > d_p + d_n$$


$$d_{3He} > d_n$$



$4qLR$

$$d_D > d_p + d_n$$

$$d_{3He} > d_n$$



$gCEDM + 4qEDM$

$$d_D \sim d_p + d_n$$

$$d_{3He} \sim d_n$$

→ $g_1 \gg g_0$, 3π -coupling (unsuppressed)

$2N$ contribution enhanced!

here: only absolute values considered


If $\bar{\theta}$ -term tests fail: effective BSM dim. 6 sources: de Vries et al. (2011)



$qEDM$

$$d_D \approx d_p + d_n$$

$$d_{3He} \approx d_n$$



$qCEDM$

$$d_D > d_p + d_n$$


$$d_{3He} > d_n$$



$4qLR$

$$d_D > d_p + d_n$$

$$d_{3He} > d_n$$



$gCEDM + 4qEDM$

$$d_D \sim d_p + d_n$$

$$d_{3He} \sim d_n$$

→ $g_1, g_0, 4N$ – coupling

$2N$ contribution difficult to assess!

here: only absolute values considered

Summary and Outlook

- θ EDM: relevant low-energy couplings **quantifiable**

strategy A: measure $d_n, d_p, d_D \xrightarrow{d_D(2N)} \bar{\theta} \xrightarrow{\text{test}} d_{3He}$

strategy A': measure $d_n, (d_p), d_{3He} \xrightarrow{d_{3He}(2N)} \bar{\theta} \xrightarrow{\text{test}} d_D$

strategy B: measure d_n (or d_p) + Lattice QCD $\rightsquigarrow \bar{\theta} \xrightarrow{\text{test}} d_D$

strategy B': measure d_n (or d_p) + Lattice QCD $\rightsquigarrow \bar{\theta} \xrightarrow{\text{test}} d_p$ (or d_n)

- qEDM, qCEDM, 4QLR:

- **NDA required** to assess sizes of low-energy couplings
- disentanglement possible by measurements of d_n, d_p, d_D & d_{3He}

- gCEDM, 4quark chiral singlet:

controlled calculation/disentanglement difficult (lattice ?)

- Ultimate progress may eventually come from Lattice QCD

↪ the \mathcal{CP} $NN\pi$ couplings may be accessible even for dim-6 sources

↪ then **quantifiable** d_D (d_{3He}) EFT predictions feasible in BSM case

Conclusions

- (Hadronic) EDMs play a key role in probing new sources of CP
- Measurements of hadronic EDMs are **low-energy measurements**
 - ↳ Predictions have to be given in the *empirical language of hadrons*
 - ↳ only reliable methods: *ChPT/EFT* and/or ultimately *Lattice QCD*
- Deuteron and helium-3 nuclei serve as isospin filters for EDMs

At least the EDMs of p , n , d , and ${}^3\text{He}$
have to be measured
to disentangle the underlying physics