

# Electric Dipole Moment Measurements at Storage Rings

**J. Pretz**

RWTH Aachen & FZ Jülich



Mainz, June 2014

# Outline


- **Introduction: Electric Dipole Moments (EDMs):**  
What is it?
- **Motivation:**  
Why is it interesting?
- **Experimental Method:**  
How to measure charged particle EDMs?
- **Results of first test measurements:**  
Spin tune and Spin Coherence time

What is it?

# Electric Dipoles

Classical definition:

$$\vec{d} = \sum_i q_i \vec{r}_i$$




# Order of magnitude

	atomic physics	hadron physics
charges	$e$	
$ \vec{r}_1 - \vec{r}_2 $	$1 \text{ \AA} = 10^{-8} \text{ cm}$	
EDM		
naive expectation	$10^{-8} e \cdot \text{cm}$	
observed	water molecule $2 \cdot 10^{-8} e \cdot \text{cm}$	

# Order of magnitude

	atomic physics	hadron physics
charges	$e$	$e$
$ \vec{r}_1 - \vec{r}_2 $	$1 \text{ \AA} = 10^{-8} \text{ cm}$	$1 \text{ fm} = 10^{-13} \text{ cm}$
EDM		
naive expectation	$10^{-8} e \cdot \text{cm}$	$10^{-13} e \cdot \text{cm}$
observed	water molecule $2 \cdot 10^{-8} e \cdot \text{cm}$	neutron $< 3 \cdot 10^{-26} e \cdot \text{cm}$

## Order of magnitude



neutron EDM of  $d_n = 3 \cdot 10^{-26} \text{ e}\cdot\text{cm}$  corresponds to separation of  $u$ – from  $d$ –quarks of  $\approx 5 \cdot 10^{-26} \text{ cm}$

## Operator $\vec{d} = q\vec{r}$

is odd under parity transformation ( $\vec{r} \rightarrow -\vec{r}$ ):

$$\mathcal{P}^{-1}\vec{d}\mathcal{P} = -\vec{d}$$

Consequences:

In a state  $|a\rangle$  of given parity the expectation value is 0:





$$\langle a|\vec{d}|a\rangle = -\langle a|\vec{d}|a\rangle$$

but if  $|a\rangle = \alpha|P = +\rangle + \beta|P = -\rangle$

in general  $\langle a|\vec{d}|a\rangle \neq 0 \Rightarrow$  i.e. molecules



# EDM of molecules



ground state: mixture of  $\Psi_s = \frac{1}{\sqrt{2}} (\Psi_1 + \Psi_2) \quad P = +$

$$\Psi_a = \frac{1}{\sqrt{2}} (\Psi_1 - \Psi_2) \quad P = -$$

(Cohen-Tannoudji, B. Diu, F. Laloë, Mécanique quantique)

## Order of magnitude

**Molecules** can have large EDM because of degenerated ground states with different parity

## Order of magnitude

**Molecules** can have large EDM because of degenerated ground states with different parity

**Elementary particles** (including hadrons) have a definite parity and cannot possess an EDM

$$P|\text{had}\rangle = \pm 1|\text{had}\rangle$$

## Order of magnitude

**Molecules** can have large EDM because of degenerated ground states with different parity

**Elementary particles** (including hadrons) have a definite parity and cannot possess an EDM

$$P|\text{had}\rangle = \pm 1|\text{had}\rangle$$

unless

$\mathcal{P}$  and time reversal  $\mathcal{T}$  invariance are violated!

# $\mathcal{T}$ and $\mathcal{P}$ violation of EDM

$\vec{d}$ : EDM


$\vec{\mu}$ : magnetic moment

both  $\parallel$  to spin

$$H = -\mu\vec{\sigma} \cdot \vec{B} - d\vec{\sigma} \cdot \vec{E}$$

$$\mathcal{T}: H = -\mu\vec{\sigma} \cdot \vec{B} + d\vec{\sigma} \cdot \vec{E}$$

$$\mathcal{P}: H = -\mu\vec{\sigma} \cdot \vec{B} + d\vec{\sigma} \cdot \vec{E}$$



$\Rightarrow$  EDM measurement tests violation of fundamental symmetries  $\mathcal{P}$  and  $\mathcal{T}(\overset{CP}{\equiv} CP)$

# Symmetries in Standard Model

	electro-mag.	weak	strong
$\mathcal{C}$	✓	✗	✓
$\mathcal{P}$	✓	✗	✓
$\mathcal{T} \xrightarrow{CPT} \mathcal{CP}$	✓	(✗)	(✓)

- $\mathcal{C}$  and  $\mathcal{P}$  are maximally violated in weak interactions (Lee, Yang, Wu)
- $\mathcal{CP}$  violation discovered in kaon decays (Cronin, Fitch) described by CKM-matrix in Standard Model
- $\mathcal{CP}$  violation allowed in strong interaction but corresponding parameter  $\theta_{QCD} \lesssim 10^{-10}$  (strong  $\mathcal{CP}$ -problem)

# Sources of $\mathcal{CP}$ –Violation

Standard Model	
<b>Weak interaction</b> CKM matrix	→ unobservably small EDMs
<b>Strong interaction</b> $\theta_{QCD}$	→ best limit from neutron EDM
beyond Standard Model	
e.g. SUSY	→ accessible by EDM measurements

Why is it interesting?



# Matter-Antimatter Asymmetry

Excess of matter in the universe:

	observed	SM prediction
$\eta = \frac{n_B - n_{\bar{B}}}{n_\gamma}$	$6 \times 10^{-10}$	$10^{-18}$


Sakharov (1967):  $\mathcal{CP}$  violation needed for baryogenesis

$\Rightarrow$  New  $\mathcal{CP}$  violating sources beyond SM needed to explain this discrepancy


They could manifest in EDMs of elementary particles

What do we know about  
EDMs?


# History of Neutron EDM



# EDM: Current Upper Limits



# EDM: Current Upper Limits




FZ Jülich: EDMs of **charged** hadrons:  $p, d, {}^3\text{He}$

# Why Charged Particle EDMs?

- no direct measurements for charged hadrons exist
- potentially higher sensitivity (compared to neutrons):
  - longer life time,
  - more stored protons/deuterons
- complementary to neutron EDM:  
 $d_d \stackrel{?}{=} d_p + d_n \Rightarrow \text{access to } \theta_{QCD}$
- EDM of one particle alone not sufficient to identify  $\mathcal{CP}$ -violating source

# Sources of $\mathcal{CP}$ Violation



How to measure charged  
particle EDMs?




# Experimental Method: Generic Idea

For **all** EDM experiments (neutron, proton, atoms, ...):

Interaction of  $\vec{d}$  with electric field  $\vec{E}$

For charged particles: apply electric field in a storage ring:



$$\frac{d\vec{s}}{dt} \propto \vec{d} \times \vec{E}$$

In general:

$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s}$$

build-up of vertical polarization  $s_{\perp} \propto |\vec{d}|$

# Experimental Requirements

- high precision storage ring  
( alignment, stability, field homogeneity)
- high intensity beams ( $N = 4 \cdot 10^{10}$  per fill)
- polarized hadron beams ( $P = 0.8$ )
- large electric fields ( $E = 10$  MV/m)
- long spin coherence time ( $\tau = 1000$  s),
- polarimetry (analyzing power  $A = 0.6$ , acc.  $f = 0.005$ )

$$\sigma_{\text{stat}} \approx \frac{1}{\sqrt{N f \tau P A E}} \Rightarrow \sigma_{\text{stat}}(1\text{year}) = 10^{-29} \text{ e}\cdot\text{cm}$$

challenge: get  $\sigma_{\text{sys}}$  to the same level

# Systematics

Major source:


Radial  $B$  field mimics an EDM effect:

- Difficulty: even small radial magnetic field,  $B_r$  can mimic EDM effect if  $:\mu B_r \approx dE_r$
- Suppose  $d = 10^{-29} \text{ e}\cdot\text{cm}$  in a field of  $E_r = 10 \text{ MV/m}$
- This corresponds to a magnetic field:

$$B_r = \frac{dE_r}{\mu_N} = \frac{10^{-22} \text{ eV}}{3.1 \cdot 10^{-8} \text{ eV/T}} \approx 3 \cdot 10^{-17} \text{ T}$$

Solution: Use two beams running clockwise and counter clockwise, separation of the two beams is sensitive to  $B_r$

## Systematics



Sensitivity needed:  $1.25 \text{ fT}/\sqrt{\text{Hz}}$  for  $d = 10^{-29} \text{ e cm}$   
(possible with SQUID technology)

# Spin Precession: Thomas-BMT Equation

$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s} = \frac{e}{m} [G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E} + \frac{m}{e s} \vec{d}(\vec{E} + \vec{v} \times \vec{B})] \times \vec{s}$$

$\Omega$ : angular precession frequency      $\vec{d}$ : electric dipole moment  
 $G$ : anomalous magnetic moment      $\gamma$ : Lorentz factor

# Spin Precession: Thomas-BMT Equation

$$\frac{d\vec{S}}{dt} = \vec{\Omega} \times \vec{S} = \frac{e}{m} [G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E} + \frac{m}{e\hbar} \vec{d}(\vec{E} + \vec{v} \times \vec{B})] \times \vec{S}$$

$\Omega$ : angular precession frequency     $\vec{d}$ : electric dipole moment

$G$ : anomalous magnetic moment     $\gamma$ : Lorentz factor

**dedicated ring:**    pure electric field,  
freeze horizontal spin motion  $\left(G - \frac{1}{\gamma^2 - 1}\right) = 0$

# Spin Precession: Thomas-BMT Equation

$$\frac{d\vec{s}}{dt} = \vec{\Omega} \times \vec{s} = \frac{e}{m} [G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E} + \frac{m}{e\hbar} \vec{d}(\vec{E} + \vec{v} \times \vec{B})] \times \vec{s}$$

$\Omega$ : angular precession frequency     $\vec{d}$ : electric dipole moment

$G$ : anomalous magnetic moment     $\gamma$ : Lorentz factor

**COSY:** pure magnetic ring  
access to EDM via motional electric field  $\vec{v} \times \vec{B}$ ,  
requires additional radio-frequency  $E$  and  $B$  fields  
to suppress  $G\vec{B}$  contribution

# Spin Precession: Thomas-BMT Equation

$$\frac{d\vec{S}}{dt} = \vec{\Omega} \times \vec{S} = \frac{e}{m} [G\vec{B} + \left(G - \frac{1}{\gamma^2 - 1}\right) \vec{v} \times \vec{E} + \frac{m}{eS} d(\vec{E} + \vec{v} \times \vec{B})] \times \vec{S}$$

$\Omega$ : angular precession frequency     $d$ : electric dipole moment

$G$ : anomalous magnetic moment     $\gamma$ : Lorentz factor

**COSY:** pure magnetic ring

access to EDM via motional electric field  $\vec{v} \times \vec{B}$ ,  
requires additional radio-frequency  $E$  and  $B$  fields  
to suppress  $G\vec{B}$  contribution


neglecting EDM term

spin tune:  $\nu_s \approx \frac{|\vec{\Omega}|}{|\omega_{cyc}|} = \gamma G, \quad (\vec{\omega}_{cyc} = \frac{e}{\gamma m} \vec{B})$



## Spin Tune $\nu_s$

Spin tune:  $\nu_s = \gamma G = \frac{\text{nb. of spin rotations}}{\text{nb. of particle revolutions}}$



deuterons:  $p_d = 1 \text{ GeV}/c$  ( $\gamma = 1.13$ ),  $G = -0.14256177(72)$

$$\Rightarrow \nu_s = \gamma G \approx -0.161$$

# Results of first test measurements

# Cooler Synchrotron COSY




COSY provides (polarized ) protons and deuterons with  
 $p = 0.3 - 3.7 \text{ GeV}/c$

⇒ **Ideal starting point for charged particle EDM searches**


# Experimental Setup

- Inject and accelerate vertically polarized deuterons to  $p \approx 1 \text{ GeV}/c$




# Experimental Setup

- Inject and accelerate vertically polarized deuterons to  $p \approx 1 \text{ GeV}/c$
- flip spin with help of solenoid into horizontal plane



# Experimental Setup

- Inject and accelerate vertically polarized deuterons to  $p \approx 1 \text{ GeV}/c$
- flip spin with help of solenoid into horizontal plane
- Extract beam slowly (in 100 s) on target
- Measure asymmetry and determine spin precession




# Asymmetry Measurements

- Detector signal  $N^{up,dn} \propto (1 \pm PA \sin(\gamma G f_{rev} t))$


$$A_{up,dn} = \frac{N^{up} - N^{dn}}{N^{up} + N^{dn}} = PA \sin(\gamma G f_{rev} t)$$

$A$ : analyzing power,  $P$ : polarization


$$A_{up,dn} = 0$$



$$A_{up,dn} = PA$$



# Polarimetry



Cross Section &  
Analyzing Power  
for deuterons

$$N_{up,dn} \propto (1 \pm P A \sin(\nu_s f_{rev} t))$$

$$A_{up,dn} = \frac{N^{up} - N^{dn}}{N^{up} + N^{dn}} = P A \sin(\nu_s f_{rev} t)$$

$A$  : analyzing power  
 $P$  : beam polarization




# Polarimeter

elastic deuteron-carbon scattering

Up/Down asymmetry  $\propto$  horizontal polarization  $\rightarrow \nu_s = \gamma G$


Left/Right asymmetry  $\propto$  vertical polarization  $\rightarrow d$



$$N_{up,dn} \propto 1 \pm PA \sin(\nu_s f_{rev} t), \quad f_{rev} \approx 781 \text{ kHz}$$


## Spin Tune $\nu_s$ measurement

- Problem: detector rate  $\approx 5$  kHz,  $f_{rev} = 781$  kHz  
 $\Rightarrow$  only 1 hit every 25th period
- not possible to use usual  $\chi^2$ -fit
- use unbinned Maximum Likelihood (under investigation)




## Spin Tune $\nu_s$ measurement

- map all events into first period ( $T = 1/(\nu_s f_{rev}) \approx 8\mu\text{s}$ ) and perform  $\chi^2$ -fit (requires knowledge of  $\nu_s f_{rev}$ )
- Analysis is done in macroscopic time bins of  $\approx 2\text{s}$




# Asymmetry in 1st period




- only works if  $T_s = \frac{1}{\nu_s f_{rev}}$  is correct.

## Scan of $\nu_s$




- set  $\nu_s = \nu_{\text{max}}$  and determine phase in macroscopic time bins of  $\approx 2\text{s}$
- allows for  $\sigma_{\nu_s} \approx 10^{-6}$


# Phase Measurements



# Phase Measurements




# Phase Measurements



- 1st derivative gives deviation from assumed spin tune



## Results: Spin Tune $\nu_s$



- Spin tune  $\nu_s$  can be determined to  $\approx 10^{-8}$  in 2 s
- Average  $\overline{\nu_s}$  in cycle ( $\approx 100$  s) determined to  $10^{-10}$   
 ( for  $G = 0$ ,  $d = 10^{-24}$  e·cm  $\Rightarrow$  spin tune =  $5 \cdot 10^{-11}$  )


## Spin Tune $\nu_s$

Experiment	Gedankenexperiment
$G \approx -0.14, d \approx 0$	$G = 0, d = 10^{-24} \text{ e cm}$
$\nu_s = \gamma G = -0.16$	$\nu_s = \frac{vm\gamma d}{es} = 5 \cdot 10^{-11}$


compare to  $\sigma(\nu_s) = 10^{-10}$  in 100 s measurement

# Results: Spin Coherence Time (SCT)

## Short Spin Coherence Time



Horizontal Asymmetry Run: 2042



unbunched beam


$$\Delta p/p = 10^{-5} \Rightarrow \Delta\gamma/\gamma = 3 \cdot 10^{-6}, T_{\text{rev}} \approx 10^{-6} \text{ s}$$

$\Rightarrow$  decoherence after  $< 1 \text{ s}$


cooled bunched beam  $\Rightarrow$  SCT  $\tau = 20 \text{ s}$

# Results: Spin Coherence Time (SCT)

## Long Spin Coherence Time




Horizontal Asymmetry Run: 2051




using correction sextupole to correct for higher order effects  
leads to SCT of  $\tau = 400$  s

# Results: Spin Coherence Time (SCT)

## Long Spin Coherence Time




Horizontal Asymmetry Run: 2051



using correction sextupole to correct for higher order effects  
leads to SCT of  $\tau = 400 \text{ s}$

# JEDI Collaboration

- **JEDI** = **J**ülich **E**lectric **D**ipole Moment **I**nvestigations
- $\approx 100$  members  
(Aachen, Dubna, Ferrara, Indiana, Ithaca, Jülich, Krakow, Michigan, Minsk, Novosibirsk, St. Petersburg, Stockholm, Tbilisi, ...)
- $\approx 10$  PhD students



# Summary & Outlook

- EDMs of elementary particles are of high interest to disentangle various sources of  $\mathcal{CP}$  violation searched for to explain matter - antimatter asymmetry in the Universe
- EDM of charged particles can be measured in storage rings

Plans in Jülich:  $10^{-24} e\text{cm}$  at COSY


$10^{-29} e\text{cm}$  with dedicated ring

- Experimentally very challenging because effect is tiny
- First promising results from test measurements at COSY

Spare



# Electron and Neutron EDM



J. M. Pendlebury &  
E.A. Hinds,  
NIMA 440(2000) 471

# EDM: SUSY Limits

## electron:

$$\text{MSSM: } \varphi \approx 1 \Rightarrow d = 10^{-24} - 10^{-27} \text{ e}\cdot\text{cm}$$

$$\varphi \approx \alpha/\pi \Rightarrow d = 10^{-26} - 10^{-30} \text{ e}\cdot\text{cm}$$

## neutron:


$$\text{MSSM: } d = 10^{-24} \text{ e}\cdot\text{cm} \cdot \sin \phi_{CP} \frac{200 \text{ GeV}}{M_{SUSY}}$$

# Electrostatic Deflectors




- Electrostatic deflectors from Fermilab ( $\pm 125\text{kV}$  at 5 cm  $\hat{=}$  5MV/m)
- large-grain Nb at plate separation of a few cm yields  $\approx$  20MV/m

# Wien Filter



Conventional design  
R. Gebel, S. Mey (FZ Jülich)




stripline design  
D. Hölscher, J. Slim  
(IHF RWTH Aachen)

# 1. Pure Magnetic Ring

$$\vec{\Omega} = \frac{e\hbar}{mc} \left( \textcolor{green}{G}\vec{B} + \frac{1}{2}\textcolor{red}{\eta}\vec{v} \times \vec{B} \right)$$

Problem:

Due to precession caused by magnetic moment, 50% of time longitudinal polarization component is  $\parallel$  to momentum, 50% of the time it is anti- $\parallel$ .




$E^*$  field in the particle rest frame  
tilts spin due to EDM up and down  
 $\Rightarrow$  **no net EDM effect**

# 1. Pure Magnetic Ring

$$\vec{\Omega} = \frac{e\hbar}{mc} \left( G\vec{B} + \frac{1}{2}\eta\vec{v} \times \vec{B} \right)$$

Problem:

Due to precession caused by magnetic moment, 50% of time longitudinal polarization component is  $\parallel$  to momentum, 50% of the time it is anti- $\parallel$ .



$E^*$  field in the particle rest frame tilts spin due to EDM up and down

$\Rightarrow$  **no net EDM effect**

Use resonant “magic Wien-Filter” in ring ( $\vec{E}_W + \vec{v} \times \vec{B}_W = 0$ ):

$E_W^* = 0 \rightarrow$  part. trajectory is not affected but

$B_W^* \neq 0 \rightarrow$  mag. mom. is influenced

$\Rightarrow$  **net EDM effect can be observed!**

## 2. Pure Electric Ring




Figure 3: An all-electric storage ring lattice for measuring the electric dipole moment of the proton. Except for having longer straight sections and separated beam channels, the all-in-one lattice of Fig. 1 is patterned after this lattice. Quadrupole and sextupole families, and tunes and lattice functions of the all-in-one lattice of Fig. 1 will be quite close to those given for this lattice in reference[3]. The match will be even closer with magnetic field set to zero for proton operation.

Brookhaven National Laboratory (BNL) Proposal

### 3. Combined $\vec{E}/\vec{B}$ ring






Figure 1: “All-In-One” lattice for measuring EDM’s of protons, deuterons, and helions.


Under discussion at Forschungszentrum Jülich (design: R. Talman)



# Summary of different options

		
1.) pure magnetic ring (Jülich)	existing (upgraded) COSY ring can be used , shorter time scale	lower sensitivity
2.) pure electric ring (BNL)	no $\vec{B}$ field needed	works only for $p$
3.) combined ring (Jülich)	works for $p, d, {}^3\text{He}, \dots$	both $\vec{E}$ and $\vec{B}$ required

# EDM Activities Around the World



# Systematics

- Splitting of beams:  $\delta y = \pm \frac{\beta c R_0 B_r}{E_r Q_y^2} = \pm 1 \cdot 10^{-12} \text{ m}$
- $Q_y \approx 0.1$ : vertical tune
- Modulate  $Q_y = Q_y^0 (1 - m \cos(\omega_m t))$ ,  $m \approx 0.1$
- Splitting causes  $B$  field of  $\approx 0.4 \cdot 10^{-3} \text{ fT}$
- in one year:  $10^4$  fills of 1000 s  $\Rightarrow \sigma_B = 0.4 \cdot 10^{-1} \text{ fT}$  per fill needed
- Need sensitivity  $1.25 \text{ fT}/\sqrt{\text{Hz}}$

# Systematics

