


## Spin-Tracking simulations in an idealized COSY model using Bmad

M. Vitz, Institut für Kernphysik 4, Forschungszentrum Jülich, 52425 Jülich, Germany,  
also at III. Physikalisches Institut B, RWTH Aachen University, 52056 Aachen, Germany  
on behalf of the JEDI Collaboration

### Motivation


- Electric Dipole Moment (EDM) is a fundamental property of a subatomic particle, similar to the Magnetic Dipole Moment (MDM).

- Source of  $\mathcal{P}$  and  $\mathcal{T}$  violation ( $\overset{CPT}{=} \mathcal{CP}$  violation) and therefore closely connected to matter antimatter asymmetry.



- EDM of charged particles can be measured in a storage ring as spin rotation is defined by EDM and MDM contribution [1].
- Vertical spin build-up is used to estimate the EDMs magnitude but also EDM-like systematic effects occur.
- Spin tracking simulations with Bmad Software Library are used to disentangle systematic effects from a real EDM signal [2].

### Invariant Spin Axis



- Expected horizontal tilt  $n_x$  due to the EDM strength  $\eta$  is described via:

$$\tan \xi_{EDM} = \frac{\eta\beta}{2G}$$

- Proof of principle was performed using the Bmad COSY model and tracking the reference particle for some thousand turns.



$$\langle \vec{n} \rangle = \frac{1}{n-1} \sum_{i=1}^{n-1} \left( \frac{\vec{s}_i \times \vec{s}_{i+1}}{|\vec{s}_i \times \vec{s}_{i+1}|} \right)$$

$$n_x = 5.61e-15 \pm 2.17e-15 \quad n_z = 3.08e-15 \pm 2.17e-15$$

$$n_y = 1.00e+00 \pm 5.09e-19$$

$$n_x = 1.61e-04 \pm 6.60e-16 \quad n_z = 2.86e-14 \pm 6.56e-16$$

$$n_y = 1.00e+00 \pm 1.80e-18$$





- Simulation result  $n_x$  in an idealized COSY model is in agreement with the expected tilt of the invariant spin axis  $\xi_{EDM}$ .
- As one cannot apply this method to an experiment a different approach to measure the EDM signal has to be used.

### Experiment at COSY

- The experiment measures the EDM using a Wien-Filter and a solenoid  $\Rightarrow$  **Resonant Wien Filter Method**
- Wien Filter gives the beam a phase dependent kick for vertical spin build-up:

$$E_x = E_0 \cdot \cos(2\pi f_{rev}|k + \nu_s| + \phi_{rel})$$



$$B_y = B_0 \cdot \cos(2\pi f_{rev}|k + \nu_s| + \phi_{rel})$$



- Amplitude of oscillation displays the EDM resonance strength [3].

$$\epsilon_{EDM} = \left( A_{WF}^2 (\phi_{WF} - \phi_{WF,0})^2 + A_{SN}^2 \left( \frac{\xi_{SN} - \xi_{SN,0}}{2 \sin(\pi \nu_{s,0})} \right)^2 \right)^{1/2} + \epsilon_0$$

- Find the fit point of **minimal resonance strength** ( $\phi_{WF,0}, \xi_{SN,0}$ )  $\Rightarrow \phi_{WF,0}$  is measured EDM plus systematic effects.



- Shift in  $\phi_{WF,0}$  is observed as soon as an EDM signal is included in the simulation  $\Rightarrow \phi_{WF,0}$  in an idealized COSY lattice with EDM signal simulated fits the expectation.

- The results of both methods are in agreement with each other  $\Rightarrow$  Systematics must be built in and be understood.

### References

- [1] T. Fukuyama and A. J. Silenko, Derivation of Generalized Thomas-Bargmann-Michel-Telegdi Equation for a Particle with Electric Dipole Moment, Int. J. Mod. Phys A28, 1350147, 2013.
- [2] D. C. Sagan, Bmad: A relativistic charged particle simulation library, Nuclear Instruments and Methods in Physics Research A, vol.558, pp.356-359, 2006.
- [3] F. Rathmann, N. N. Nikolaev and J. Slim, Spin dynamics investigations for the electric dipole moment experiment, Physical Review Accelerators and Beams 23, 024601, 2020.